

# Gunnpod RX

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

**WARNING:** Important safety information is provided in a boxed warning in the [full CMI](#). Read before using this medicine.

### 1. Why am I using Gunnpod RX?

Gunnpod RX contains the active ingredient nicotine. Gunnpod RX is an alternative source of nicotine to help reduce the withdrawal symptoms associated with nicotine dependency. Gunnpod RX may aid in smoking cessation by replacing the nicotine you are used to getting from cigarettes.

For more information, see Section [1. Why am I using Gunnpod RX?](#) in the full CMI.

### 2. What should I know before I use Gunnpod RX?

Do not use if you have ever had an allergic reaction to nicotine or any of the ingredients listed at the end of the CMI.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see Section [2. What should I know before I use Gunnpod RX?](#) in the full CMI.

### 3. What if I am taking other medicines?

Some medicines may interfere with Gunnpod RX and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### 4. How do I use Gunnpod RX?

- Follow all directions given to you by your doctor carefully. They may differ from the information contained within this leaflet.
- Follow the instructions provided and use Gunnpod RX until your doctor tells you to stop.
- Do not exceed the recommended dosage.

More instructions can be found in Section [4. How do I use Gunnpod RX?](#) in the full CMI.

### 5. What should I know while using Gunnpod RX?

|                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Things you should do</b>        | <ul style="list-style-type: none"><li>• Follow the directions given to you by your doctor carefully.</li><li>• Remind any doctor, dentist or pharmacist you visit that you are using nicotine.</li><li>• Check with your doctor if you are pregnant or intend to become pregnant.</li><li>• Tell your doctor if you have allergies to any other medication.</li></ul>                                                                                                                                                                                           |
| <b>Things you should not do</b>    | <ul style="list-style-type: none"><li>• Do not stop using this medicine suddenly – you may experience nicotine withdrawal symptoms.</li><li>• Do not use this medicine if you are allergic to nicotine, or any of the ingredients listed at the end of this leaflet.</li><li>• Do not use this medicine if you are a non-smoker or do not have an existing nicotine dependence.</li><li>• Do not use this medicine if you are under 18 years or age.</li><li>• Do not give this medicine to anyone else, even if they have the same condition as you.</li></ul> |
| <b>Driving or using machines</b>   | <ul style="list-style-type: none"><li>• Be careful before you drive or use and machines or tools until you know how Gunnpod RX affects you. Gunnpod RX may cause dizziness in some people.</li></ul>                                                                                                                                                                                                                                                                                                                                                            |
| <b>Looking after your medicine</b> | <ul style="list-style-type: none"><li>• Keep Gunnpod RX pod sealed in the pack until needed and store in a cool dry place below 25°C, away from moisture heat or sunlight.</li><li>• Store Gunnpod RX where young children cannot reach it.</li></ul>                                                                                                                                                                                                                                                                                                           |

For more information, see Section [5. What should I know while using Gunnpod RX?](#) in the full CMI.

## 6. Are there any side effects?

If you smoke or use any other nicotine-container product while you are using Gunnpod RX, you may suffer an overdose of nicotine. However, if used correctly, overdose is unlikely. A large overdose may present symptoms such as exhaustion, convulsions, low blood pressure, circulatory collapse and breathing difficulties. In this event, you should immediately phone the Poisons Information Centre (by calling 13 11 26), or contact your doctor, or go to the Emergency Department at your nearest hospital.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

# Gunnpod RX

**Active ingredient(s):** *Nicotine* (ni-kuh-teen)

## Consumer Medicine Information (CMI)

This leaflet provides important information about using Gunnpod RX. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Gunnpod RX<sup>1</sup>.**

**Where to find information in this leaflet:**

- [1. Why am I using Gunnpod RX?](#)
- [2. What should I know before I use Gunnpod RX?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Gunnpod RX?](#)
- [5. What should I know while using Gunnpod RX?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

## 1. Why am I using Gunnpod RX?

**Gunnpod RX contains the active ingredient nicotine.**

Gunnpod RX is an alternative source of nicotine to help reduce the withdrawal symptoms associated with nicotine dependency. Gunnpod RX may aid in smoking cessation by replacing the nicotine you are used to getting from cigarettes.

**Gunnpod RX is used to aid in smoking cessation.**

Gunnpod RX pod contains a dilute nicotine liquid. When used in conjunction with the device the liquid is vaporised for inhalation which replaces the nicotine you would otherwise get from smoking. The device and the cartridge when used together provides nicotine to relieve some of the unpleasant symptoms such as feeling ill or irritable that smokers often feel when they try to give up. The nicotine from Gunnpod RX can also reduce your cravings for a cigarette and help you to resist the urge to smoke.

This type of treatment through inhalation is a type Nicotine Replacement Therapy (NRT) to aid in smoking cessation.

## 2. What should I know before I use Gunnpod RX?

### Warnings

**Do not use Gunnpod RX if:**

- you are allergic to nicotine, or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine.
- you are a non-smoker.
- you don't have an existing nicotine dependence.
- you are under 18 years of age.

**Check with your doctor if you:**

- have any other medical conditions such as:
  - high blood pressure
  - problems with blood circulation
  - a history of chest pain, heart attack or stroke
  - blood vessel disorder like Buerger's disease or Raynaud's syndrome
  - liver or kidney disease
  - stomach or duodenal ulcers
  - adrenal gland tumour
  - insulin dependent diabetes
  - an overactive thyroid
  - asthma or other breathing disorders
- take any medicines for any other condition
- have allergies to any other medication

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

### Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

Smoking during pregnancy has risk such as poor growth of your baby before birth, premature birth or still birth.

Stopping smoking is the best way to improve both your health and that of your baby. The earlier you stop smoking the better.

**Gunnpod RX should not be used while breastfeeding.**

### 3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Gunnpod RX.**

### 4. How do I use Gunnpod RX?

#### How much to use

- Follow all directions given to you by your doctor carefully. They may differ from the information contained within this leaflet.
- If you do not understand the instructions, ask your doctor or pharmacist for help.
- Follow the instructions provided and use Gunnpod RX until your doctor tells you to stop.

#### When to use Gunnpod RX

- Gunnpod RX should be used when you feel the need to smoke a cigarette. It should be used as an alternative to smoking.

#### How to use Gunnpod RX

- Remove Gunnpod RX pod from the sealed packaging
- Take off the mouthpiece cap of the pod
- Insert pod into GPRX 650 Device
- Place your lips around the mouthpiece and slowly inhale.
- The blue light at the bottom will light up when inhaling. If the blue light is blinking, it indicates low battery. If there is no blue light on the Gunnpod RX, the battery has run out – cease use immediately and dispose of the product.
- Inhale the vapour into your lungs (Do not swallow it).
- Exhale.
- You will know that Gunnpod RX is empty when there is insufficient liquid in the vape or the battery has run out. Once Gunnpod RX is empty, cease use immediately and dispose of the Gunnpod RX safely.
- Using a cotton swab with isopropyl alcohol, clean the mouthpiece and bottom air intake outlet and then dry thoroughly. This will prevent any residue build-up.
- Keep your Gunnpod RX pod sealed in its pack until required for use.
- Always ensure you carefully follow all directions given to you by your prescriber. Always enquire with your doctor or pharmacist if you are unclear about or do not understand how to use your device.

#### If you forget to use Gunnpod RX

Gunnpod RX is intended for use as a replacement for cigarette smoking to aid in smoking cessation. Always use

Gunnpod RX in line with the directions of your doctor or pharmacist.

If you stop using Gunnpod RX abruptly, you may experience symptoms of nicotine withdrawal. Common symptoms of nicotine withdrawal include:

- Feeling tense and irritable
- Depression
- Appetite changes
- Constipation and gas
- Insomnia
- Difficulty concentrating
- Cough and cold-like symptoms

#### If you use too much Gunnpod RX

If you think that you have used too much Gunnpod RX, you may need urgent medical attention.

If you smoke or use any other nicotine containing product while you are using Gunnpod RX, you may suffer an overdose of nicotine. However, if used correctly, nicotine overdose is unlikely. Symptoms of nicotine overdose include:

- Headache
- Dizziness
- Upset stomach
- Drooling
- Vomiting
- Diarrhoea
- Cold sweat
- Blurred vision
- Hearing distortion
- Confusion
- Weakness

#### You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

**You should do this even if there are no signs of discomfort or poisoning.**

### 5. What should I know while using Gunnpod RX?

#### Things you should do

- Follow the directions given to you by your doctor carefully.
- Check with your doctor if you are pregnant or intend to become pregnant.
- Tell your doctor if you have allergies to any other medication.

#### Call your doctor straight away if you:

- Are pregnant or become pregnant while using Gunnpod RX.

- Experience any side effects while using Gunnpod RX.
- Notice anything else that may be making you feel unwell.

Remind any doctor, dentist or pharmacist you visit that you are using Gunnpod RX.

### Things you should not do

- Do not stop using this medicine suddenly – you may experience nicotine withdrawal symptoms, making you feel unwell.
- Do not use this medicine if you are allergic to nicotine, or any of the ingredients listed at the end of this leaflet.
- Do not use this medicine if you are a non-smoker or do not have an existing nicotine dependence, as this medicine is to aid smoking cessation.
- Do not use this medicine if you are under 18 years of age, this medicine should be kept out of reach of children.
- Do not give your medicine to anyone else, even if they have the same condition as you.

### Using Gunnpod RX in conjunction with cigarette smoking is discouraged.

Gunnpod RX is intended to aid smoking cessation by replacing cigarettes.

### Driving or using machines

#### Be careful before you drive or use any machines or tools until you know how Gunnpod RX affects you.

Gunnpod RX may cause dizziness in some people.

### Looking after your medicine

- Keep Gunnpod RX pod sealed in the blister pack until needed.
- Store in a cool dry place below 25°C.

Store it away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on windowsills.

Keep it where young children cannot reach it.

### When to discard your medicine

Do not use this medicine after the expiry date (EXP) printed on the pack. If you use it after the expiry date has passed, it may not work as well.

Do not use if the packaging is torn, shows signs of tampering, or if the blister pack is damaged.

### Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

## 6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

Nicotine vaping products have the potential to aid with smoking cessation efforts, but they may have unwanted side effects in a few people. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some side effects.

Stopping smoking itself can cause some symptoms such as dizziness, headache, sleep disturbance, cough and cold-like symptoms. Symptoms such as depression, irritability, anxiety and insomnia may also be related to withdrawal symptoms associated with giving up smoking.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Do not be alarmed by the following list of side effects. You may not experience any of them.

### Less serious side effects

| Less serious side effects                                                                                                                                                                                                                                                                                                                                                                                                                                                          | What to do                                                                                                              |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>• Nausea</li> <li>• Upset stomach</li> <li>• Vomiting</li> <li>• Diarrhoea</li> <li>• Dizziness</li> <li>• Headache</li> <li>• Dry mouth</li> <li>• Throat irritation</li> <li>• Indigestion</li> <li>• Constipation</li> <li>• Sleep disturbances</li> <li>• Tremors</li> <li>• Nervousness</li> <li>• Tiredness or weakness</li> <li>• Increased sweating</li> <li>• Exacerbation of asthma in people with pre-existing asthma</li> </ul> | <p><b>Speak to your doctor or pharmacist if you have any of these less serious side effects and they worry you.</b></p> |

### Serious side effects

| Serious side effects                                                                                                                                                                                                         | What to do                                                                                                                             |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>• Fast or irregular heartbeat (tachycardia)</li> <li>• Swelling of your face, lips mouth or throat</li> <li>• Difficulty breathing/shortness of breath.</li> <li>• Seizure</li> </ul> | <p><b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of</b></p> |

|  |                                    |
|--|------------------------------------|
|  | <b>these serious side effects.</b> |
|--|------------------------------------|

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

**Reporting side effects**

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

**7. Product details**

This medicine is only available with a doctor's prescription.

**What Gunnpod RX contains**

|                                                     |                                                                        |
|-----------------------------------------------------|------------------------------------------------------------------------|
| <b>Active ingredient<br/>(main ingredient)</b>      | Nicotine<br>(equivalent to 20mg/mL)                                    |
| <b>Other ingredients<br/>(inactive ingredients)</b> | Glycerol<br>Propylene Glycol<br>Water<br>Menthol Flavour<br>Stabiliser |

**Do not take this medicine if you are allergic to any of these ingredients.**

**What Gunnpod RX looks like**

Gunnpod RX is available in 1 flavour:

- Menthol

Gunnpod RX is available in 20 mg/mL (100 mg total nicotine per cartridge).

Actual usage may vary depending on individual usage patterns, draw duration and device maintenance

Gunnpod RX is not on the Australian Register of Therapeutic Goods (ARTG) and is referred to as an unapproved therapeutic good.

Unapproved therapeutic goods have not been evaluated by the Therapeutic Goods Administration (TGA) for their quality, safety or effectiveness.

**Who distributes Gunnpod RX**

Entity C Pty Ltd  
Suite 353/ 585 Little Collins Street,  
Melbourne VIC 3000

This leaflet was prepared in Nov 2025.